

DOLORCLAST RADIAL SHOCKWAVE THERAPY

There is a new form of treatment available at Coburg Osteopathy. Dolorclast Radial Shockwave Therapy is useful for the treatment of pain related to a number of chronic soft tissue conditions including;

- Tennis/golfers elbow
- Plantar fasciitis
- Achilles tendinopathy
- Patellar tendinopathy
- Shin splints
- Bursitis
- ITB syndrome
- Rotator cuff tendonitis
- Hamstring tendonitis
- Chronic trigger points

This form of treatment was developed from other medical indications (treating kidney stones) and is widely used in Europe and North America for treatment of the above soft tissue conditions. There are numerous published scientific studies, which you are free to examine, the majority of which demonstrate positive outcomes. Most importantly it has been demonstrated as a safe treatment option. Radial shockwaves are high energy soundwaves transmitted from a probe held against and passing through the skin and spreading outwards into the underlying tissues. This is believed to induce increased blood flow and metabolic activity around the site of pain accelerating the healing process. At the very least it effectively reduces pain from this area.

ABOUT THE TREATMENT:

After a simple examination of the affected area, shockwaves are applied via a handpiece or applicator held against the skin. The initial phase of treatment may cause some deep pain however this indicates correct targeting of the problem area. This is usually followed by numbness or heaviness in the area and the latter phase of the treatment feels less painful. The treatment sessions are of 10 minutes duration during which you may receive up to 2000 pulses. Many patients get pain relief just days after the first treatment. Depending on the condition and your response you may require from 1 to 5 treatments, although 3 is usually sufficient.

AFTER THE TREATMENT:

You may feel some soreness which may intensify on the night of treatment. Simple analgesia and icing may be required to control this pain. There may also be some minor bruising in the treatment area. You should rest from aggravating activities for 2-3 days after treatment as your doctor will discuss.

SIDE EFFECTS:

Post-treatment pain as above.

Local bruising – uncommon and minor.

Tendon rupture has not been reported following radial shockwave therapy, unlike cortisone injection and surgery.

CONTRAINDICATIONS:

Please inform your doctor if you have a one of the following contra-indications as this form of treatment may not be appropriate for you –

- Pregnancy
- Bleeding and blood coagulation disorders and associated medication consumption.
- Skin wound or acute inflammation in treatment area.
- Tumour in treatment area.

COST:

\$95 per treatment session

RESULTS:

Obviously the human body is a complex structure and like many things in medicine we can provide no guarantees of effectiveness of treatment. We can only keep up-to-date with the latest quality scientific research and make this available to you to make your decision.

If you have any questions please feel free to discuss these with your doctor.